

The Area of Polynomial Images and Preimages

Edward Crane *

Department of Pure Mathematics and Statistics, University of Cambridge,
Cambridge CB3 0WA, United Kingdom.

February 1, 2008

Abstract

Let p be a monic complex polynomial of degree n and let K a measurable subset of the complex plane. We show that the area of $p(K)$, counted with multiplicity, is at least $\pi n (\text{Area}(K)/\pi)^n$ and that

$$\frac{\text{Area}(p^{-1}(K))}{\pi} \leq \left(\frac{\text{Area}(K)}{\pi} \right)^{1/n}.$$

Both bounds are sharp. The special case of the latter result in which K a disc was proved by Pólya in 1928. We use Carleman's isoperimetric inequality relating the conductance and area for plane condensers. We include a summary of the necessary potential theory.

1 Introduction and Statement of Results

When p is a polynomial of degree n over $\mathbb{C}$, the set

$$E(p, r) = \{z \in \mathbb{C} : |p(z)| = r^n\}$$

is called a lemniscate, after Bernoulli's lemniscate $\{z \in \mathbb{C} : |z^2 - 1| = 1\}$. It is natural to ask how large the set enclosed by a lemniscate can be. We need to make some normalisation for this to be meaningful; the simplest is to ask for p to be monic. Perhaps surprisingly, the area of the lemniscate $E(p, 1)$ is then bounded independently of p . In fact the following sharp inequality was proved by Pólya in 1928, [5].

*Supported by an EPSRC research studentship

Theorem 1 (Pólya's inequality).

Let p be a monic polynomial of degree n over $\mathbb{C}$ and let D be a disc in $\mathbb{C}$. Then the Euclidean area of $p^{-1}(D)$ is at most $\pi \left(\frac{\text{Area}(D)}{\pi} \right)^{1/n}$, with equality only when $p : z \mapsto a(z - b)^n + c$ and the center of D is c , the unique critical value of p .

It is natural to ask whether we can get a larger preimage by fixing the area of D but allowing its shape to vary. The main theorem of this paper is that we cannot.

Theorem 2. *Let p be a monic polynomial of degree n over $\mathbb{C}$. Let K be any measurable subset of the plane. Then*

$$\text{Area}(p^{-1}(K)) \leq \pi \left(\frac{\text{Area}(K)}{\pi} \right)^{1/n},$$

with equality if and only if K is (up to sets of measure zero) a disc and p has a unique critical value at the center of that disc.

Theorem 2 is a simple consequence of the following stronger theorem:

Theorem 3. *Let p be a monic polynomial of degree n over $\mathbb{C}$, and K be any measurable subset of the plane. Define the multiplicity $n(z, p, K)$ to be the number of p -preimages of z in K , counted according to their valency. Then the area of $p(K)$ counted with multiplicity satisfies*

$$\int_{\mathbb{C}} n(z, p, K) dA(z) = \int_K |p'(z)|^2 dA \geq n\pi \left(\frac{\text{Area}(K)}{\pi} \right)^n,$$

with equality if and only if K is (up to sets of measure zero) a disc and p has a unique critical value at the center of that disc.

For a compact set $K \subset \mathbb{C}$, define

$$\rho(K) = \frac{\text{Area}(K)}{\pi \text{cap}(K)^2},$$

where $\text{cap}(K)$ is the logarithmic capacity of K . Then $\rho(K)$ is a measure of the *roundness* of K : we will see that $\rho(K) \in [0, 1]$, and $\rho(K) = 1$ if and only if K is a full-measure subset of a disc. We use ρ to formulate the following scale-invariant version of theorem 2.

Theorem 4. *If p is any complex polynomial of degree n , not necessarily monic, and K is any compact subset of the plane, then*

$$\rho(p^{-1}(K)) \leq \rho(K)^{1/n}.$$

This corollary is sharp for each value of ρ . To see this we can take K to be the union of the unit disc with a radial line segment, and $p : z \mapsto z^n$, so that we still get equality in theorem 2.

In section 2 we give a quick introduction to the potential theory that we will need. In section 3 we discuss some isoperimetric inequalities and their relationship with Pólya's inequality (theorem 1), which we prove since it is an important ingredient in the proof of theorem 2. Theorems 2, 3 and 4 are proved in section 4.

A survey of area estimates for lemniscates has recently been given by Lubinsky [4], with a view towards applications in the convergence theory of Padé approximation. In [2], Eremenko and Hayman made progress on the related problem of bounding the length of $E(p, r)$. Fryntov and Rossi [3] have obtained the sharp analogue of Pólya's inequality (theorem 1) bounding the hyperbolic area of the preimage of a hyperbolic disc under a finite Blaschke product. This raises the question of finding the sharp Blaschke product analogues of theorems 2 and 3.

The author thanks Assaf Naor and Ben Green for posing the question that led to this paper, and his PhD supervisor Keith Carne for useful conversations.

2 Capacity of plane subsets and condensers

Definition 1. : *A plane condenser is a pair (E, B) of subsets of $\mathbb{C}$, where $E \neq \mathbb{C}$ is open and B is a non-empty closed subset of E .*

The terminology arises from the fact that a pair of conducting cylinders with cross-section ∂E and B respectively could be used as a condenser (or capacitor). The *capacity* of the condenser (E, B) is physically the capacitance per unit length of an infinitely long pair of such cylinders. The same quantity describes the *conductance* between ∂E and B of an isotropic resistor consisting of a plate in the shape of $E \setminus B$. We compute this conductance by considering the electrical potential f that would be induced in

U if we were to connect ∂E to an electrical potential 0 and B to potential +1. Given f , we can compute the current $-\nabla f$ that would flow in response to the potential f , and the power consumed is proportional to

$$L(f) = \int_{E \setminus B} |\nabla f|^2 dA,$$

We expect a physical potential f to minimise $L(f)$ over all possible potential functions satisfying the given boundary conditions. Calculus of variations tells us that if there is an extremal f , it must be harmonic on $E \setminus B$. We call such an f a Green's function for the condenser. A Green's function only exists if the boundary $\partial E \cup \partial B$ is regular for the Dirichlet problem, but we avoid this difficulty by defining

$$\text{cap}(E, B) = \frac{1}{4\pi} \inf L(f),$$

where the infimum is taken over all continuously differentiable $f : \mathbb{C} \rightarrow \mathbb{R}$ such that $f = 0$ on $\mathbb{C} \setminus E$ and $f = 1$ on B . We call such functions *admissible* for the condenser (E, B) . Note that $\text{cap}(E, B)$ may be zero, as it is when B is a finite set. From the definition it is immediate that capacity is monotonic, i.e.

$$E \subseteq F \text{ and } B \supseteq C \implies \text{cap}(E, B) \geq \text{cap}(F, C).$$

Lemma 1. *Suppose that on some open set $U \subset \mathbb{C}$ we have an analytic function ψ such that each point of E has exactly n preimages in U , counted according to valency. Then*

$$\text{cap}(\psi^{-1}(E), \psi^{-1}(B)) = n \text{cap}(E, B).$$

Proof. Suppose that $f : \mathbb{C} \rightarrow \mathbb{R}$ is any admissible function for (E, B) . The hypothesis implies that the restriction of ψ to $\psi^{-1}(E)$ is a proper map, so we can extend $f \circ \psi$ to get an admissible function for $(\psi^{-1}(E), \psi^{-1}(B))$ by giving it the value 0 outside U . Since ψ is almost everywhere conformal,

$$\begin{aligned} L(f \circ \psi) &= \int_{\psi^{-1}(E \setminus B)} |\nabla(f \circ \psi)(z)|^2 d\text{Area}(z) \\ &= \int_{\psi^{-1}(E \setminus B)} |(\nabla f)(\psi(z))| \cdot |\psi'(z)|^2 d\text{Area}(z) \\ &= n \int_{E \setminus B} |\nabla f(w)|^2 d\text{Area}(w) = n L(f). \end{aligned}$$

□

In particular the capacity is a *conformal invariant* of condensers: if $\varphi : E \rightarrow \mathbb{C}$ is a univalent function then

$$\text{cap}(E, B) = \text{cap}(\varphi(E), \varphi(B)).$$

For example, if $(E \setminus B)$ is a ring domain then its modulus is $1/(4\pi \text{cap}(E, B))$.

Let K be a compact set in the plane. A Green's function for K is a continuous function $f : \mathbb{C} \rightarrow \mathbb{R}$, zero on K and harmonic on $\mathbb{C} \setminus K$, with $f(z) = \log |z| - \log t + o(1)$ as $z \rightarrow \infty$. If K has a Green's function then the logarithmic capacity of K is defined to be $\text{cap}(K) = t$. For general K we define $\text{cap}(K) = \inf \text{cap}(J)$ over all compact sets $J \supset K$ with regular boundary for the Dirichlet problem on $\mathbb{C} \setminus J$. By pulling back Green's functions, it is easy to verify that if p is a monic polynomial of degree n then

$$\text{cap}(p^{-1}(B)) = \text{cap}(B)^{1/n}.$$

3 Isoperimetric Inequalities

A relationship between capacity and 2-dimensional Lebesgue measure is given by the following 'isoperimetric' inequality:

Theorem 5. (Carleman, 1918)

$$\frac{1}{\text{cap}(E, B)} \leq \log \left(\frac{\text{Area}(E)}{\text{Area}(B)} \right),$$

with equality iff E and B are concentric discs.

The proof of Carleman's inequality uses the fact that the Dirichlet integral $L(f)$ does not increase when f is replaced by its Schwarz symmetrization, the function $S(f)$ whose superlevel sets are concentric discs with the same area as the corresponding level sets of f . For details, see the classic book of Pólya and Szegő, [6], or [1] for a more modern account.

Taking $E = B(0, R)$ and then letting $R \rightarrow \infty$ in Carleman's isoperimetric inequality yields the following well-known isoperimetric theorem for logarithmic capacity. For a simple proof, including the equality case, see theorem 5.3.5 in [7].

Theorem 6. For any compact set $K \subset \mathbb{C}$,

$$\text{Area}(K) \leq \pi \text{cap}(K)^2,$$

with equality if and only if K is a disc.

We have now collected everything we need to prove Pólya's inequality, theorem 1. The capacity of the disc D is precisely the radius of D , so

$$\begin{aligned}\text{cap}(D) &= \left(\frac{\text{Area}(D)}{\pi} \right)^{1/2}, \\ \text{cap}(p^{-1}(D)) &= \left(\frac{\text{Area}(D)}{\pi} \right)^{1/2n},\end{aligned}$$

and, applying theorem 6,

$$\text{Area}(p^{-1}(D)) \leq \pi \left(\frac{\text{Area}(D)}{\pi} \right)^{1/n},$$

as required. In view of the strong link between logarithmic capacity and polynomials, theorems 1 and 6 are virtually equivalent. In [4], Pólya's inequality is proved using Gronwall's area formula, and used to deduce the isoperimetric inequality for logarithmic capacity.

4 Proof of theorems 2 and 3

Lemma 2. *For any complex polynomial g of degree d ,*

$$\int_{\mathbb{C}} |g(w)| \mathbf{1}_{|g(w)| \leq x} dA \geq \frac{2x}{d+2} \text{Area}(\{w \in \mathbb{C} : |g(w)| \leq x\}).$$

Proof. By lemma 1, we have

$$\text{cap}(g^{-1}(B(0, x)), g^{-1}(B(0, s))) = \frac{d}{2(\log x - \log s)}.$$

Theorem 5 gives

$$\frac{\text{Area}(\{w \in \mathbb{C} : s \leq |g(w)| \leq x\})}{\text{Area}(\{w \in \mathbb{C} : |g(w)| \leq x\})} \geq 1 - \left(\frac{s}{x}\right)^{2/d},$$

so

$$\begin{aligned}\int_{\mathbb{C}} |g(w)| \mathbf{1}_{|g(w)| \leq x} dA &= \int_0^x \text{Area}(\{w \in \mathbb{C} : s \leq |g(w)| \leq x\}) ds \\ &\geq \text{Area}(\{w \in \mathbb{C} : |g(w)| \leq x\}) \int_0^x 1 - \left(\frac{s}{x}\right)^{2/d} ds \\ &= \frac{2x}{d+2} \text{Area}(\{w \in \mathbb{C} : |g(w)| \leq x\}).\end{aligned}$$

□

Now fix a monic polynomial p and $A > 0$. Among all measurable sets K with $\text{Area}(K) = A$, the Dirichlet integral

$$\int_K |p'(w)|^2 d\text{Area}(w)$$

is minimised when K is the sublevel set

$$K_t = \{w \in \mathbb{C} : |p'(w)|^2 \leq t\}.$$

Here t is determined uniquely by the condition $\text{Area}(K_t) = A$. The polynomial $z \mapsto (p'(z)/n)^2$ is monic, with degree $2n - 2$, so theorem 1 gives

$$A = \text{Area}(K_t) \leq \pi \left(\frac{\pi(t/n^2)^2}{\pi} \right)^{1/(2n-2)}.$$

Rearranging this we have

$$t \geq n^2 \left(\frac{A}{\pi} \right)^{n-1}.$$

Now we apply lemma 2 to the polynomial $g = (p')^2$ to obtain

$$\begin{aligned} \int_{K_t} |p'(w)|^2 dA(w) &= \int_{\mathbb{C}} |p'(w)|^2 \mathbf{1}_{|p'(w)|^2 \leq t} dA(z) \\ &\geq \frac{2t}{2n} \text{Area}(K_t) = \frac{tA}{n} \\ &\geq n\pi \left(\frac{A}{\pi} \right)^n. \end{aligned}$$

For equality, we must have equality in our application of Pólya's inequality, so p must be $p : z \mapsto (z - b)^n + c$, and K can differ from disc K_t at most by a set of 2-dimensional Lebesgue measure zero. This completes the proof of theorem 3.

To obtain theorem 2, observe that a monic polynomial p maps $p^{-1}(K)$ onto K with multiplicity n everywhere, so

$$\text{Area}(K) = \frac{1}{n} \int_{p^{-1}(K)} |p'(w)|^2 dA(w).$$

Finally, theorem 4 is obtained by dividing both sides of the inequality of theorem 2 by $\text{cap}(K)^2 = \text{cap}(p^{-1}(K))^2$.

References

- [1] Bandle, Catherine. *Isoperimetric Inequalities and Applications*, Pitman, 1980.
- [2] Alexandre Eremenko and Walter Hayman, *On the length of lemniscates*, Michigan J. Math **46** (1999), 409–415.
- [3] Alexander Fryntov and John Rossi, *Hyperbolic symmetrization and an inequality of Dyn'kin*, Entire functions in modern analysis (Tel–Aviv, 1997), Israel Math. Conf. Proc., **15**, 103–115.
- [4] Doron Lubinsky, *Small values of polynomials: Cartan, Pólya and others*, Journal of Inequalities and Applications, **1**, (1997), no. 3, 199–222.
- [5] G. Pólya, *Beitrag zur Verallgemeinerung des Verzerrungssatzes auf mehrfach zusammenhängende Gebiete*, S-B. Akad. Wiss, Berlin (1928), 228–232 & 280–282. Also in volume 1 of Pólya's Collected Papers, MIT press, (1974).
- [6] G. Pólya and G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*, Annals of Mathematics Studies **27**, Princeton University Press, (1951).
- [7] Thomas Ransford, *Potential Theory in the Complex Plane*, London Mathematical Society Student Texts **28**, Cambridge University Press (1995).